

PATENT
Attorney Docket No. 033275-083

BE IT KNOWN, that I, MATTHEW C. COFFEY, a resident of Calgary, Canada, have
invented new and useful improvements in

**METHODS FOR THE TREATMENT OF CELLULAR
PROLIFERATIVE DISORDERS**

BURNS, DOANE, SWECKER & MATHIS, LLP.

METHODS FOR THE TREATMENT OF CELLULAR PROLIFERATIVE DISORDERS

Related Application

This application claims the benefit of U.S. Provisional Patent Application
5 Ser. No. 60/246,728, filed November 9, 2000, the disclosure of which is hereby
incorporated by reference in its entirety.

Field of the Invention

The present invention relates to methods of identifying the susceptibility of
cells to reovirus infection by measuring constitutive ras-MAP signaling. The
10 invention also pertains to methods using reovirus for the treatment of cellular
proliferative disorders, and particularly cellular proliferative disorders wherein the
proliferating cells exhibit constitutive MAPK phosphorylation, in mammals. In
particular, the methods provide for reovirus treatment of mammals to treat
proliferative disorders which include breast tumors, a subset of tumors in which
15 mutation of the ras gene is not believed to play a significant role.

References

The following publications, patent applications and patents are cited in this
application:

U.S. Patent No. 5,023,252.

20 WO 99/08692, published February 25, 1999.

Archer *et al.* (1995), *Br. J. Cancer* 72:1259-1266.

Armstrong, G.D. *et al.* (1984), *Virology* 138:37.

Barbacid, M., *Annu. Rev. Biochem.*, 56:779-827 (1987).

Baselga *et al.* (1996), *J. Clin. Onc.* 14:737-744.

- Bos, J. (1989) *Cancer Res.* 49:4682.
- Carter *et al.* (1992), *PNAS* 89:4285-4289.
- Chandron and Nibert, "Protease cleavage of reovirus capsid protein mu1 and mu1C is blocked by alkyl sulfate detergents, yielding a new type of infectious subvirion particle", *J. of Virology* 72(1):467-75 (1998).
- 5 Chaubert, P. *et al.* (1994), *Am. J. Path.* 144:767.
- Clark *et al.* (1995), *Science* 268:233-239.
- Clark *et al.* (1996), *Intl. J. Cancer* 65:186-191.
- Cuff *et al.*, "Enteric reovirus infection as a probe to study immunotoxicity of the gastrointestinal tract" *Toxicological Sciences* 42(2):99-108 (1998)
- 10 DiDomenico *et al.* (1996), *Cancer Res.* 56:4516-4521
- Duncan *et al.*, "Conformational and functional analysis of the C-terminal globular head of the reovirus cell attachment protein" *Virology* 182(2):810-9 (1991).
- Dvorak *et al.* (1988), *Am J Path* 133:95-109.
- 15 Fields, B.N. *et al.* (1996), *Fundamental Virology, 3rd Edition*, Lippincott-Raven.
- Gentsch, J.R.K. and Pacitti, A.F. (1985), *J. Virol.* 56:356.
- E. Harlow and D. Lane, "Antibodies: A laboratory manual", Cold Spring Harbor Laboratory (1988).
- Harweth *et al.* (1992), *J Biol Chem* 267:15160-15167.
- 20 Hudziak *et al.* (1989), *Mol Cell Biol* 9:1165-1172.
- Hung *et al.* (1995), *Gene* 159:65-71.
- Jacobs *et al.* (1983), *Cancer Res* 43:1696-1702.
- Janes, P.W., *et al.* (1994) *Oncogene* 9:3601.
- Jardines *et al.* (1993) *Pathobiology* 61:268-282.
- 25 Koenders *et al.* (1991), *Cancer Res* 51:4544-4548.

- Lee. J.M. *et al.* (1993) *PNAS* 90:5742-5746.
- Lee, P.W.K. *et al.* (1981) *Virology*, 108:134-146.
- Levitzki, A. (1994) *Eur. J. Biochem.* 226:1.
- Lowe. S.W. *et al.* (1994) *Science*, 266:807-810.
- 5 Mah *et al.*, "The N-terminal quarter of reovirus cell attachment protein sigma 1 possesses intrinsic virion-anchoring function" *Virology* 179(1):95-103 (1990).
- Migliaccio *et al.* (1996), *EMBO J* 15:1292-1300.
- Migliaccio *et al.* (1998), *EMBO J* 17:2008-2018.
- Millis, NE *et al.* (1995) *Cancer Res.* 55:1444.
- 10 Nagy *et al.* (1989) *Biochim Biophys Acta* 948:305-326.
- Paul R.W. *et al.* (1989) *Virology* 172:382-385.
- Pietras *et al.* (1994), *Oncogene* 9:1829-1838.
- Raybaud-Diogene. H. *et al.* (1997) *J. Clin. Oncology*, 15(3):1030-1038.
- Remington's *Pharmaceutical Sciences*, Mace Publishing Company, Philadelphia
15 PA 17th ed. (1985).
- Rosen, L. (1960) *Am. J. Hyg.* 71:242.
- Sabin, A.B. (1959), *Science* 130:966.
- Shackney *et al.* (1998), *Clin Cancer Res* 4:913-928.
- Slamon *et al.* (1989), *Science* 244:707-712.
- 20 Smith, R.E. *et al.*, (1969) *Virology*, 39:791-800.
- Spandidos (1987), *Anticancer Res* 7:991-996.
- Stanley, N.F. (1967) *Br. Med. Bull.* 23:150.
- Strong, J.E. *et al.*, (1993) *Virology*, 197:405-411.
- Strong, J.E. and Lee, P.W.K., (1996) *J. Virol.*, 70:612-616.

Strong, J.E. *et al.*, (1998) *EMBO J*, 17:3351-3362.

Turner and Duncan, "Site directed mutagenesis of the C-terminal portion of reovirus protein signal: evidence for a conformation-dependent receptor binding domain" *Virology* 186(1):219-27 (1992).

5 Verbeek *et al.* (1996) *J Path* 180:383-388.

Wiessmuller, L. and Wittinghofer, F. (1994), *Cellular Signaling* 6(3):247-267.

Zhou *et al.* (1989), *Oncogene* 4:105-108.

All of the above publications, patent applications and patents are herein incorporated by reference in their entirety to the same extent as if each individual
10 publication, patent application or patent was specifically and individually indicated to be incorporated by reference in its entirety.

State of the Art

Normal cell proliferation is regulated by a balance between growth-promoting proto-oncogenes and growth-constraining tumor-suppressor genes.
15 Tumorigenesis can be caused by genetic alterations to the genome that result in the mutation of those cellular elements that govern the interpretation of cellular signals, such as potentiation of proto-oncogene activity or inactivation of tumor suppression. It is believed that the interpretation of these signals ultimately influences the growth and differentiation of a cell, and that misinterpretation of
20 these signals can result in neoplastic growth (neoplasia).

Genetic alteration of the proto-oncogene Ras is believed to contribute to approximately 30% of all human tumors (Wiessmuller, L. and Wittinghofer, F. (1994), *Cellular Signaling* 6(3):247-267; Barbacid, M. (1987) *A Rev. Biochem.*
25 56, 779-827). The role that Ras plays in the pathogenesis of human tumors is specific to the type of tumor. Activating mutations in Ras itself are found in most types of human malignancies, and are highly represented in pancreatic cancer (80%), sporadic colorectal carcinomas (40-50%), human lung adenocarcinomas

(15-24%), thyroid tumors (50%) and myeloid leukemia (30%) (Millis, NE *et al.* (1995) *Cancer Res.* 55:1444; Chaubert, P. *et al.* (1994), *Am. J. Path.* 144:767; Bos, J. (1989) *Cancer Res.* 49:4682). Ras activation is also demonstrated by upstream mitogenic signaling elements, notably by tyrosine receptor kinases
5 (RTKs). These upstream elements, if amplified or overexpressed, ultimately result in elevated Ras activity by the signal transduction activity of Ras. Examples of this include overexpression of PDGFR in certain forms of glioblastomas, as well as in c-erbB-2/neu in breast cancer (Levitzki, A. (1994) *Eur. J. Biochem.* 226:1; Janes, P.W., *et al.* (1994) *Oncogene* 9:3601; Bos, J. (1989) *Cancer Res.*
10 49:4682).

Current methods of treatment for neoplasia include surgery, chemotherapy and radiation. Surgery is typically used as the primary treatment for early stages of cancer; however, many tumors cannot be completely removed by surgical means. In addition, metastatic growth of neoplasms may prevent complete cure of
15 cancer by surgery. Chemotherapy involves administration of compounds having antitumor activity, such as alkylating agents, antimetabolites, and antitumor antibiotics. The efficacy of chemotherapy is often limited by severe side effects, including nausea and vomiting, bone marrow depression, renal damage, and central nervous system depression. Radiation therapy relies on the greater ability
20 of normal cells, in contrast with neoplastic cells, to repair themselves after treatment with radiation. Radiotherapy cannot be used to treat many neoplasms, however, because of the sensitivity of tissue surrounding the tumor. In addition, certain tumors have demonstrated resistance to radiotherapy and such may be dependent on oncogene or anti-oncogene status of the cell (Lee, J.M. *et al.* (1993)
25 *PNAS* 90:5742-5746; Lowe, S.W. *et al.* (1994) *Science*, 266:807-810; Raybaud-Diogene, H. *et al.* (1997) *J. Clin. Oncology*, 15(3):1030-1038).

Breast cancer is amongst the most common and feared cancer in women with an estimated one in eight American women developing it in their lifetime (NCI-SEER, 1998). Although there have been tremendous achievements in breast

cancer treatment that have allowed a stable mortality rate in the face of an increasing incident rate, most of these advances can be attributed to improved early *detection* methods and not novel *treatment* strategies. The fact that there have been few advances in the actual treatment of this affliction demands that the
5 development of unconventional treatment strategies be explored.

Never in the history of cancer biology has there been a more promising period in the development of therapeutics for the treatment of breast cancer. Many of these therapeutics are targeted at specific receptors that are overexpressed in subsets of this cancer, and are now finding some clinical use. Perhaps one of the
10 most promising of these new therapies is the use of antibodies targeted to the HER2 receptor that is frequently overexpressed in breast cancer. Results from the use of these monoclonal antibodies directed at the extracellular moieties of this growth factor receptor are promising, and indeed have met with some clinical success (Baselga, 1996; for reviews see Nass, 1998; Hung, 1995). There are
15 however a number of technical obstacles that must be addressed. Firstly, these monoclonal antibodies directed against HER2 are not themselves cytocidal. but are rather cytostatic (Pietras, 1994; Harwerth, 1992; Hudziak, 1989; Carter, 1992). Importantly, in a SCID or nude mouse model, anti-HER2 antibodies are effective only in *repressing* tumor growth and not actually causing tumor *regression*.
20 Further, this repression of tumor growth is alleviated rapidly once the antibody administration is ceased. On a more positive note, there is evidence that treatment of tumors with anti-HER2 antibodies in an immune competent animal may actually result in a cytocidal effect mediated by antibody-dependent cellular cytotoxicity (ADCC) (Carter, 1992). Secondly, targeting growth factor receptor
25 (EGFR) may be of limited value, as not all breast cancers express these receptors at elevated levels. Additionally, this strategy targets only receptor tyrosine kinases and would have little therapeutic benefit for those breast cancers that overexpress non-receptor tyrosine kinases such as Src, which have already been implicated in the growth of some breast cancers (Clark, 1996). Finally, the biology of the
30 tumor blood vessels must also be taken into account as far as the capacities to

deliver these large macromolecules to their target cells. It has been demonstrated that the vasculature of a tumor is heterogeneous and the ability of a vessel to leak large macromolecules, such as antibodies, is a function of their spatial orientation to the tumor mass (Dvorak, 1988; Nagy and Dvorak, 1989). It has been shown
5 that the vessels with the greatest permeability to these molecules reside primarily at the tumor-host interface, and the least permeable vessels are those that actually penetrate the tumour mass (Dvorak, 1988). The result of this differential permeability is that these tumour specific antibodies, and their cytotoxic effector cells, would fail to penetrate into the tumour mass and would be only effective on
10 the periphery of the mass.

Despite the fact that activating mutations in *ras* are infrequent in breast cancer, there is a growing body of evidence that activation of the Ras/MAPK pathway is important in the initiation and progression of this disease. Upstream elements of the Ras pathway, notably receptor tyrosine kinases (RTK) are
15 frequently overexpressed in breast cancers. HER-2/neu (erbB-2) is overexpressed in approximately 30% of all breast cancers (Spandidos 1987; Zhou, 1989; Archer, 1995) and is associated with poor patient prognosis (Slamon, 1989). When overexpressed in NIH-3T3 cells, HER-2 mediates transformation; however, there does appear to be a threshold level of overexpression that is required for this
20 transformation to occur (Jardines, 1993; Clark 1995). It appears that this transforming ability of HER-2 is dependent upon Ras activity as cell lines overexpressing HER-2 exhibit a dramatic increase in MAP kinase activity, the latter being reflective of Ras activity (Janes, 1994). The growth factor receptor EGFR that is closely related to, but distinct from HER-2, has also been observed
25 to be overexpressed in breast tumors and has also been correlated strongly with a poor patient prognosis (Shackney, 1998; Koenders, 1991).

Additional elements upstream of Ras have also been implicated in the etiology of breast cancer. The non-receptor tyrosine kinase c-Src has been implicated as being as important candidate in promoting the progression of breast

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235

a U87 tumor xenograft and following palpable tumor establishment the tumors were treated with a single intratumor injection of reovirus. This single treatment resulted in dramatic tumor regression.

In view of the drawbacks associated with the current means for treating neoplastic growth, the need still exists for improved methods for the treatment of most types of cancers and, particularly, breast cancer. It would be useful to have a means available to determine reovirus susceptibility to predict the effectiveness of reovirus treatment.

SUMMARY OF THE INVENTION

The present invention is directed to methods of identifying the susceptibility of cells to reovirus infection by measuring constitutive ras-MAP signaling. The invention also provides methods using reovirus for the treatment of cellular proliferative disorders, and particularly cellular proliferative disorders wherein the proliferating cells exhibit constitutive MAPK phosphorylation, in mammals. In particular, the methods provide for reovirus treatment of mammals to treat proliferative disorders which include breast tumors, a subset of tumors in which mutation of the ras gene is not believed to play a significant role.

This invention also pertains to methods for preventing reovirus recognition in the treatment of cellular proliferative disorders, and particularly cellular proliferative disorders which exhibit constitutive MAPK phosphorylation, in mammals. The mammal may be selected from dogs, cats, sheep, goats, cattle, horses, pigs, mice, humans and non-human primates. The method comprises administering to the proliferating cells an effective amount of one or more reoviruses under conditions which result in substantial lysis of the proliferating cells.

The reovirus may be a mammalian reovirus or an avian reovirus. The reovirus may be modified such that the outer capsid is removed, the virion is packaged in a liposome or micelle or the proteins of the outer capsid have been mutated. The reovirus agent can be administered in a single dose or in multiple
5 doses. The proliferative disorder may be a neoplasm. Both solid and hematopoietic neoplasms can be targeted. Use of preceding, concurrent or subsequent immunosuppression may result in more effective reovirus treatment.

Accordingly, in one aspect the invention provides a method of determining cellular susceptibility to reovirus infection by measuring constitutive ras-MAP
10 signaling, wherein the presence of said signaling indicates susceptibility to infection by reovirus. Constitutive ras-MAP signaling results in activation of the MAP kinase whether there is a mitogen or not, and activation of the MAP kinase leads to phosphorylation of MAP kinase. Therefore, the state of MAP kinase phosphorylation can be determined as a measure of constitutive ras-MAP
15 signaling. The state of MAP kinase phosphorylation can be determined by any method established in the art, and in particular by using an antibody specific for phosphorylated MAP kinase.

Accordingly, the present invention can be used to diagnose proliferative disorders which can be treated with reovirus. Thus, a biological sample can be
20 collected from a mammal suspected of having a proliferative disorder and cells in the sample are tested for constitutive ras-MAP signaling according to the present invention. The proliferative disorder may be any condition associated with abnormally active cell growth, such as neurofibromatosis. More preferably, the proliferative disorder is selected from the group consisting of lung cancer, prostate
25 cancer, colorectal cancer, thyroid cancer, renal cancer, adrenal cancer, liver cancer, pancreatic cancer, breast cancer and central and peripheral nervous system cancer. A particularly preferred proliferative disorder is breast cancer.

The present invention is applicable to any animal with a proliferative disorder. Preferably, the animal is a mammal. More preferably, the animal is selected from the group consisting of dogs, cats, sheep, goats, cattle, horses, pigs, mice, non-human primates, and humans. Most preferably, the animal is human.

5 In a further aspect, the invention provides a method of treating a proliferative disorder in a mammal, which disorder is characterized by proliferating cells which exhibit constitutive MAPK phosphorylation, comprising administering to the proliferating cells in said mammal an effective amount of one or more reoviruses under conditions which result in substantial lysis of the
10 proliferating cells. Methods which further comprise a step selected from the group consisting of: administering to the proliferating cells in said mammal an effective amount of an immune suppressive agent; removing B-cells or T-cells from said mammal; removing anti-reovirus antibodies from said mammal; removing antibodies from said mammal; administering anti-antireovirus antibodies to said
15 mammal; and suppressing the immune system of the mammal are also provided, as are methods which further comprise administration of a chemotherapeutic agent.

Also provided is a method of treating a neoplasm in a human, which neoplasm is characterized by proliferating cells which exhibit constitutive MAPK phosphorylation, comprising administering to the neoplasm a reovirus in an
20 amount sufficient to result in substantial oncolysis of the neoplastic cells. Preferably, the reovirus is administered systemically or by injection into or near a solid neoplasm. Methods further comprising the step of previously, concurrently or subsequently suppressing or otherwise compromising the immune system of the mammal are also included.

25 Also provided is a method of inhibiting metastasis of a neoplasm in a mammal, which neoplasm is characterized by proliferating cells which exhibit constitutive MAPK phosphorylation, comprising administering to the mammal a reovirus in an amount sufficient to result in substantial lysis of the neoplastic cells.

Methods further comprising the step of previously, concurrently or subsequently suppressing or otherwise compromising the immune system of the mammal are also included.

Also provided is a method of treating a suspected neoplasm in a mammal,
5 which neoplasm is characterized by proliferating cells which exhibit constitutive MAPK phosphorylation, comprising surgical removal of substantially all of the neoplasm and administration of an effective amount of reovirus at or near to the surgical site resulting in oncolysis of any remaining neoplastic cells. The reovirus may also be administered systemically. Methods further comprising the step of
10 previously, concurrently or subsequently suppressing or otherwise compromising the immune system of the mammal are also included.

Also provided is a pharmaceutical composition comprising an effective amount of a reovirus and a pharmaceutically acceptable excipient. Also provided is a pharmaceutical composition comprising an immunosuppressant or
15 immunoinhibitant, a reovirus and a pharmaceutically acceptable excipient. Kits comprising a reovirus and, optionally, an immunosuppressant or immunoinhibitant are also provided.

The methods and pharmaceutical compositions of the invention provide an effective means to treat neoplasia, without the side effects associated with other
20 forms of cancer therapy. When used, inhibition or suppression of the immune system increases the availability of reovirus to infect and lyse proliferating cells which exhibit constitutive MAPK phosphorylation because anti-reovirus antibodies are not formed. Because reovirus is not known to be associated with disease, any safety concerns associated with deliberate administration of a virus are minimized.

25 The foregoing and other objects, features and advantages of the invention will be apparent from the following more particular description of preferred embodiments of the invention, as illustrated in the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the effect of *v-src* transformation on host cell susceptibility to reovirus infection. Uninfectable NIH-3T3 parental cells and *v-src* transformed NIH-3T3 cells were grown to 80% confluency in a 24 well plate and were then
5 exposed to reovirus at an estimated MOI of 80 PFU per cell. Cells and media were harvested at the indicated times post infection and resultant lysate was used for plaque titration assay. *V-src* transformed NIH-3T3 cells (closed circles), NIH-3T3 parentals (open circles). (Mean \pm standard deviation).

Figures 2A to 2C illustrate *in vitro* reovirus replication in human breast
10 tumor cell lines. (A) Reovirus protein synthesis in mock-infected and reovirus-infected breast cancer cell lines. Cells were labeled with [³⁵S]-methionine from 46 to 48 hours after infection. Lysates were prepared and subsequently immunoprecipitated with a polyclonal anti-reovirus type 3 serum and then analyzed by SDS-PAGE. Reovirus proteins are indicated on the right. (B) Reovirus
15 infectivity correlates with constitutive MAPK phosphorylation. The breast tissue cell line HBL-100 and the breast tumor cell lines MDA-MB-468, MCF7, MDA-MB-435, T47D and SK-BR-3 were plated in a six well plate. Cells were either grown in the presence of 10% FCS or were serum starved (0.5% FCS) for a 48 hour period. Monolayers were washed in PBS and cell lysates were prepared and
20 subjected to SDS-PAGE. Following blotting onto nitrocellulose paper, samples were probed with antibodies directed against phospho-MAPK. Phospho-MAPK levels were all standardized by total MAPK levels. (C) Total MAPK levels in the human breast tissue and breast cancer cell lines.

Figure 3 illustrates reovirus-mediated *in vivo* oncolysis against human
25 breast tumor xenograft. SCID mice were implanted subcutaneously and unilaterally with MDA-MB-468 human breast tumor xenografts. Following the establishment of a palpable tumor mass, a single intratumoral injection of either

live reovirus (open circles) or UV-inactivated virus (closed circles) was administered and tumor growth was followed for a period of four weeks. (Mean \pm SEM).

DETAILED DESCRIPTION OF THE INVENTION

5 Previously we have shown that the human reovirus could be used as an effective oncolytic agent against human glioblastoma xenografts in a SCID mouse model (WO 99/08692). Here we have found that cellular susceptibility to reovirus infection may be determined by measuring constitutive ras-MAP signaling of the cell, since the presence of such signaling indicates susceptibility to infection
10 by reovirus. We also present evidence that reovirus is useful as an oncolytic agent against mammary tumors. Although *ras* mutations are infrequent in the etiology of breast cancer, aberrant Ras/MAPK signaling via upstream signal elements, such as receptor tyrosine kinases and non-receptor tyrosine kinases (such as c-Src), is common.

15 For the same reasons that the treatment of breast cancers with monoclonal antibodies may be impaired (see discussion above), reovirus may be an attractive cancer therapy. Firstly, reovirus itself is a cytocidal agent and does not rely upon immune effector cells to cause tumor regression. Indeed, the natural mechanism of killing infected cells is via direct lysis due to viral replication (Tyler and Fields,
20 1996). Secondly, it targets those breast cancers in which there is activation of Ras. This activation is not restricted to activating mutations of Ras (admittedly a rare subset of breast tumors) but also includes activation of Ras caused by elements upstream of Ras itself. These elements include not only receptor tyrosine kinases such as EGFR and HER2, but also include non-receptor tyrosine kinases
25 such as the Src family members. Taken together, this type of therapy could be used with great efficacy against a tumor type as heterogeneous as breast cancer and would not be as restricted a strategy as one that targets only on receptor. Finally,

the inability of antibodies to penetrate into solid tumor masses suggests that reovirus, if delivered intratumorally, should replicate unheeded. Thus, it is useful to know whether the cells of a proliferative disorder are susceptible to infection by reovirus in order to predict the efficacy of such treatment.

5 To assess whether Src family kinases could mediate reovirus infection, uninfected NIH-3T3 cells were transformed with *v-src* and were then challenged with reovirus. We determined that *v-src* signaling confers infectability to reovirus infection.

 We next examined five breast cancer cell lines; MDA-MB-468, MCF7,
10 MDA-MB-435, T47D, and SK-BR-3 as well as a cell line derived from normal breast tissue, HBL-100 for *in vitro* reovirus replication. All five of the tumor derived cell lines were infectable by reovirus while the HBL-100 was unable to effectively replicate the virus.

 To determine if the Ras pathway was indeed activated in these cell lines,
15 the level of MAPK phosphorylation was assessed in the presence and absence of serum. Those cell lines that were infectable exhibited constitutive MAPK phosphorylation even in the absence of mitogen, while the cell line which was not susceptible to reovirus infection displayed MAPK phosphorylation only in the presence of serum. Therefore, constitutive MAPK phosphorylation is an
20 indication of susceptibility to reovirus infection.

 To determine if reovirus could be used as an oncolytic agent *in vivo* against breast tumors, SCID mice were implanted with MDA-MB-468 human tumour xenographs. Following the establishment of palpable tumours, the mice were treated with a single injection of reovirus and tumour size was monitored for a
25 four-week period. The single injection resulted in dramatic regression of the tumour size. Finally, the ability of reovirus to act against primary breast cancer tumours was determined, and the results indicate that reovirus was capable of

replicating in biopsy samples collected from various patients. Therefore, many breast tumors are susceptible to reovirus infection.

The name reovirus (Respiratory and enteric orphan virus) is a descriptive acronym suggesting that these viruses, although not associated with any known disease state in humans, can be isolated from both the respiratory and enteric tracts (Sabin, A.B. (1959), *Science* 130:966). The term "reovirus" refers to all viruses classified in the reovirus genus.

Reoviruses are viruses with a double-stranded, segmented RNA genome. The virions measure 60-80 nm in diameter and possess two concentric capsid shells, each of which is icosahedral. The genome consists of double-stranded RNA in 10-12 discrete segments with a total genome size of 16-27 kbp. The individual RNA segments vary in size. Three distinct but related types of reovirus have been recovered from many species. All three types share a common complement-fixing antigen.

The human reovirus consists of three serotypes: type 1 (strain Lang or T1L), type 2 (strain Jones, T2J) and type 3 (strain Dearing or strain Abney, T3D). The three serotypes are easily identifiable on the basis of neutralization and hemagglutinin-inhibition assays (Sabin, A.B. (1959), *Science* 130:966; Fields, B.N. *et al.* (1996), Fundamental Virology, 3rd Edition, Lippincott-Raven; Rosen, L. (1960) *Am. J. Hyg.* 71:242; Stanley, N.F. (1967) *Br. Med. Bull.* 23:150).

Although reovirus is not known to be associated with any particular disease, many people have been exposed to reovirus by the time they reach adulthood (i.e., fewer than 25% in children < 5 years old, to greater than 50% in those 20-30 years old (Jackson G.G. and Muldoon R.L. (1973) *J. Infect. Dis.* 128:811; Stanley N.F. (1974) *In: Comparative Diagnosis of Viral Diseases*, edited by E. Kurstak and K. Kurstak, 385-421, Academic Press, New York).

For mammalian reoviruses, the cell surface recognition signal is sialic acid (Armstrong, G.D. *et al.* (1984), *Virology* 138:37; Gentsch, J.R.K. and Pacitti, A.F. (1985), *J. Virol.* 56:356; Paul R.W. *et al.* (1989) *Virology* 172:382-385) Due to the ubiquitous nature of sialic acid, reovirus binds efficiently to a multitude
5 of cell lines and as such can potentially target many different tissues; however, there are significant differences in susceptibility to reovirus infection between cell lines.

As described herein, Applicant have discovered that cells which exhibit constitutive MAPK phosphorylation are susceptible to reovirus infection.
10 "Resistance" of cells to reovirus infection indicates that infection of the cells with the virus did not result in significant viral production or yield. Cells that are "susceptible" are those that demonstrate induction of cytopathic effects, viral protein synthesis, and/or virus production. Resistance to reovirus infection was found to be at the level of gene translation, rather than at early transcription:
15 while viral transcripts were produced, virus proteins were not expressed.

The implantation of human tumor cells into SCID mice is recognized as a well known model system for testing the effectiveness of various anti-tumor agents in humans. It has previously been shown that pharmaceuticals effective against human tumors implanted into SCID mice are predictive of their effectiveness
20 against the same tumors in humans.

Based upon these discoveries, Applicant have developed methods for determining susceptibility to reovirus infection by measuring constitutive ras-MAP signaling and methods for treating proliferative disorders in mammals wherein the proliferating cells exhibit constitutive MAPK phosphorylation. Representative
25 mammals include dogs, cats, sheep, goats, cattle, horses, pigs, mice, non-human primates, and humans. In a preferred embodiment, the mammal is a human.

In the diagnostic methods of the invention, the level of MAPK phosphorylation of proliferating cells is determined in the presence or absence of mitogen. The presence of such constitutive ras-MAP signaling in the cells is indicative of susceptibility to reovirus infection.

5 In the treatment methods of the invention, reovirus is administered to proliferating cells in the individual mammal exhibiting constitutive MAPK phosphorylation. In one embodiment of this invention a course of reovirus therapy is administered one or more times.

10 In the methods of the invention, reovirus is administered to proliferating cells in the individual mammal exhibiting constitutive MAPK phosphorylation. Representative types of human reovirus that can be used include type 1 (e.g., strain Lang or T1L); type 2 (e.g., strain Jones or T2J); and type 3 (e.g., strain Dearing or strain Abney, T3D or T3A); other strains of reovirus can also be used. In a preferred embodiment, the reovirus is human reovirus serotype 3, more
15 preferably the reovirus is human reovirus serotype 3, strain Dearing. Alternatively, the reovirus can be a non-human mammalian reovirus (e.g., non-human primate reovirus, such as baboon reovirus; equine; or canine reovirus), or a non-mammalian reovirus (e.g., avian reovirus). A combination of different serotypes and/or different strains of reovirus, such as reovirus from different
20 species of animal, can be used.

The reovirus may be naturally occurring or modified. The reovirus is "naturally-occurring": when it can be isolated from a source in nature and has not been intentionally modified by humans in the laboratory. For example, the reovirus can be from a "field source": for example, from a patient.

25 The reovirus may be modified but still capable of lytically infecting a mammalian cell exhibiting constitutive MAPK phosphorylation. The reovirus may be chemically or biochemically pretreated (e.g., by treatment with a protease, such

as chymotrypsin or trypsin) prior to administration to the proliferating cells. Pretreatment with a protease removes the outer coat or capsid of the virus and may increase the infectivity of the virus. The reovirus may be coated in a liposome or micelle (Chandron and Nibert, "Protease cleavage of reovirus capsid protein mu1 and mu1C is blocked by alkyl sulfate detergents, yielding a new type of infectious subviral particle", *J. of Virology* 72(1):467-75 (1998)) to reduce or prevent an immune response from a mammal which has developed immunity to the reovirus. For example, the virion may be treated with chymotrypsin in the presence of micelle forming concentrations of alkyl sulfate detergents to generate a new infectious subviral particle (ISVP). An ISVP may be used either alone or in combination with whole virus to provide an agent that is either poorly recognized or has not been previously prevented by the patient's immune system.

The reovirus may be a recombinant reovirus from two or more types of reoviruses with differing pathogenic phenotypes such that it contains different antigenic determinants thereby reducing or preventing an immune response by a mammal previously exposed to a reovirus subtype. Such recombinant virions, also known as reassortants, can be generated by co-infection of mammalian cells with different subtypes of reovirus with the resulting resorting and incorporation of different subtype coat proteins into the resulting virion capsids.

The reovirus may be modified by incorporation of mutated coat proteins, such as for example $\sigma 1$, into the virion outer capsid. The proteins may be mutated by replacement, insertion or deletion. Replacement includes the insertion of different amino acids in place of the native amino acids. Insertions include the insertion of additional amino acid residues into the protein at one or more locations. Deletions include deletions of one or more amino acid residues in the protein. Such mutations may be generated by methods known in the art. For example, oligonucleotide site directed mutagenesis of the gene encoding for one of the coat proteins could result in the generation of the desired mutant coat protein. Expression of the mutated protein in reovirus infected mammalian cells *in vitro*

such as COS1 cells will result in the incorporation of the mutated protein into the reovirus virion particle (Turner and Duncan, "Site directed mutagenesis of the C-terminal portion of reovirus protein sigma1: evidence for a conformation-dependent receptor binding domain" *Virology* 186(1):219-27 (1992); Duncan et al., "Conformational and functional analysis of the C-terminal globular head of the reovirus cell attachment protein" *Virology* 182(2):810-9 (1991); Mah et al., "The N-terminal quarter of reovirus cell attachment protein sigma 1 possesses intrinsic virion-anchoring function" *Virology* 179(1):95-103 (1990))

The reovirus is preferably a reovirus modified to reduce or eliminate an immune reaction to the reovirus. Such modified reovirus are termed "immunoprotected reovirus". Such modifications could include packaging of the reovirus in a liposome, a micelle or other vehicle to mask the reovirus from the mammals immune system. Alternatively, the outer capsid of the reovirus virion particle may be removed since the proteins present in the outer capsid are the major determinant of the host humoral and cellular responses.

A "proliferative disorder" is any cellular disorder in which the cells proliferate more rapidly than normal tissue growth. Thus a "proliferating cell" is a cell that is proliferating more rapidly than normal cells. The proliferative disorder, includes but is not limited to neoplasms. A neoplasm is an abnormal tissue growth, generally forming a distinct mass, that grows by cellular proliferation more rapidly than normal tissue growth. Neoplasms show partial or total lack of structural organization and functional coordination with normal tissue. These can be broadly classified into three major types. Malignant neoplasms arising from epithelial structures are called carcinomas, malignant neoplasms that originate from connective tissues such as muscle, cartilage, fat or bone are called sarcomas and malignant tumors affecting hematopoietic structures (structures pertaining to the formation of blood cells) including components of the immune system, are called leukemias and lymphomas. A tumor is the neoplastic growth of the disease cancer. As used herein, a "neoplasm", also referred to as a "tumor",

is intended to encompass hematopoietic neoplasms as well as solid neoplasms. Other proliferative disorders include, but are not limited to neurofibromatosis.

At least some of the cells of the proliferative disorder which are treated using the methods of the present invention exhibit constitutive MAPK

- 5 phosphorylation. Similarly, at least some of the cells which are determined to be susceptible to reovirus infection using the methods of the present invention exhibit constitutive MAPK phosphorylation.

"B-cells" refers to B-lymphocytes. There are two major subpopulations of B lymphocytes, B-1 and B-2 cells. B-1 cells are self-renewing and frequently
10 secrete high levels of antibody which binds to a range of antigens (polyspecificity) with a relatively low affinity. The majority of B cells, B-2 cells, are directly generated from precursors in the bone marrow and secrete highly specific antibody.

"T-cells" refers to T-lymphocytes. T-cells differentiate within the thymus
15 gland and are specialized to operate against cells bearing intracellular organisms. T-cells only recognize antigen when it is on the surface of a body cell.

"Anti-reovirus antibody" refers to an antibody which binds to reovirus. "IgG antibodies" refers to immunoglobulin G antibodies. IgG, the most abundant type of antibody, carries the major burden of neutralizing bacterial toxins and
20 binding to microorganisms to enhance their phagocytosis. "Humanized antibodies" refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Administration to a proliferating cell or neoplasm" indicates that the
25 reovirus is administered in a manner so that it contacts the proliferating cells or cells of the neoplasm (also referred to herein as "neoplastic cells"). The route by

which the reovirus is administered, as well as the formulation, carrier or vehicle, will depend on the location as well as the type of the neoplasm. A wide variety of administration routes can be employed. For example, for a solid neoplasm that is accessible, the reovirus can be administered by injection directly to the neoplasm.

- 5 For a hematopoietic neoplasm, for example, the reovirus can be administered intravenously or intravascularly. For neoplasms that are not easily accessible within the body, such as metastases or brain tumors, the reovirus is administered in a manner such that it can be transported systemically through the body of the mammal and thereby reach the neoplasm (e.g., intrathecally, intravenously or
10 intramuscularly). Alternatively, the reovirus can be administered directly to a single solid neoplasm, where it then is carried systemically through the body to metastases. The reovirus can also be administered subcutaneously, intraperitoneally, topically (e.g., for melanoma), orally (e.g., for oral or esophageal neoplasm), rectally (e.g., for colorectal neoplasm), vaginally (e.g., for
15 cervical or vaginal neoplasm), nasally or by inhalation spray (e.g., for lung neoplasm).

- Reovirus can be administered systemically to mammals which are immune compromised or which have not developed immunity to the reovirus epitopes. In such cases, reovirus administered systemically, i.e. by intravenous injection, will
20 contact the proliferating cells resulting in lysis of the cells. Where the mammals to be treated have higher titers of anti-reovirus antibodies, more reovirus must be administered to be effective.

- Immunocompetent mammals previously exposed to a reovirus subtype may have developed humoral and/or cellular immunity to that reovirus subtype.
25 Nevertheless, it has been found that direct injection of the reovirus into a solid tumor in immunocompetent mammals will result in the lysis of the neoplastic cells. On the other hand, when the reovirus is administered systemically to immunocompetent mammals, the mammals may produce an immune response to the reovirus. Such an immune response may be avoided if the reovirus is of a

subtype to which the mammal has not developed immunity, or the reovirus has been modified as previously described herein such that it is immunoprotected, for example, by protease digestion of the outer capsid or packaging in a micelle.

Alternatively, it is contemplated that the immunocompetency of the mammal against the reovirus may be suppressed either by the prior or co-administration of pharmaceuticals known in the art to suppress the immune system in general (Cuff et al., "Enteric reovirus infection as a probe to study immunotoxicity of the gastrointestinal tract" *Toxicological Sciences* 42(2):99-108 (1998)) or alternatively the administration of such immunoinhibitors as anti-antireovirus antibodies. The humoral immunity of the mammal against reovirus may also be temporarily reduced or suppressed by plasmaphoresis of the mammals blood to remove the anti-reovirus antibodies. The humoral immunity of the mammal against reovirus may additionally be temporarily reduced or suppressed by the intravenous administration of non-specific immunoglobulin to the mammal.

It is contemplated that the reovirus may be administered to immunocompetent mammals immunized against the reovirus in conjunction with the administration of immunosuppressants and/or immunoinhibitors. Such immunosuppressants and immunoinhibitors are known to those of skill in the art and include such agents as cyclosporin, rapamycin, tacrolimus, mycophenolic acid, azathioprine and their analogs, and the like. Other agents are known to have immunosuppressant properties as well (see, e.g., Goodman and Gilman, 7th Edition, page 1242, the disclosure of which is incorporated herein by reference). Such immunoinhibitors also include "anti-antireovirus antibodies," which are antibodies directed against anti-reovirus antibodies. Such antibodies can be made by methods known in the art. See for example "Antibodies: A laboratory manual" E. Harlow and D. Lane, Cold Spring Harbor Laboratory (1988). Such anti-antireovirus antibodies may be administered prior to, at the same time or shortly after the administration of the reovirus. Preferably an effective amount of the anti-antireovirus antibodies are administered in sufficient time to reduce or eliminate an

immune response by the mammal to the administered reovirus. The terms
"immunosuppressant" or "immune suppressive agent" include conventional
immunosuppressants, immunoinhibitors, antibodies, and conditions such as
radiation therapy or HIV infection which result in compromise of the immune
5 system.

The term "substantial lysis" means at least 10% of the proliferating cells
are lysed, more preferably of at least 50% and most preferably of at least 75% of
the cells are lysed. The percentage of lysis can be determined for tumor cells by
measuring the reduction in the size of the tumor in the mammal or the lysis of the
10 tumor cells *in vitro*.

A "mammal suspected of having a proliferative disorder" means that the
mammal may have a proliferative disorder or tumor or has been diagnosed with a
proliferative disorder or tumor or has been previously diagnosed with a
proliferative disorder or tumor, the tumor or substantially all of the tumor has been
15 surgically removed and the mammal is suspected of harboring some residual tumor
cells.

This invention also includes pharmaceutical compositions which contain, as
the active ingredient, one or more immunosuppressants or immunoinhibitors and
one or more of the reoviruses associated with "pharmaceutically acceptable
20 carriers or excipients". In making the compositions of this invention, the active
ingredients/ immunosuppressant or immunoinhibitor and reovirus are usually
mixed with an excipient, diluted by an excipient or enclosed within such a carrier
which can be in the form of a capsule, sachet, paper or other container. When the
pharmaceutically acceptable excipient serves as a diluent, it can be a solid, semi-
25 solid, or liquid material, which acts as a vehicle, carrier or medium for the active
ingredient. Thus, the compositions can be in the form of tablets, pills, powders,
lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups,
aerosols (as a solid or in a liquid medium), ointments containing, for example, up

to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, sterile water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

For preparing solid compositions such as tablets, the principal active ingredients/ immunosuppressant or immunoinhibitor and reovirus are mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules.

The tablets or pills of the present invention may be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids

and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous
5 solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as corn oil, cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or
10 mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described herein. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably pharmaceutically acceptable solvents may be nebulized by use of inert gases. Nebulized solutions may be inhaled
15 directly from the nebulizing device or the nebulizing device may be attached to a face mask tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

Another preferred formulation employed in the methods of the present
20 invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the reovirus of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, for example, U.S. Patent 5,023,252, herein incorporated by
25 reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Other suitable formulations for use in the present invention can be found in *Remington's Pharmaceutical Sciences*, the disclosure of which is incorporated herein by reference.

5 The immunosuppressant or immunoinhibitor and reovirus or the pharmaceutical composition comprising the immunosuppressant or immunoinhibitor and reovirus may be packaged into convenient kits providing the necessary materials packaged into suitable containers. It is contemplated the kits may also include chemotherapeutic agent.

10 The immunosuppressant or immunoinhibitor is administered in an appropriate amount and using an appropriate schedule of administration sufficient to result in immunosuppression or immunoinhibition of the mammal's immune system. Such amounts and schedules are well known to those of skill in the art.

15 The reovirus is administered in an amount that is sufficient to treat the proliferative disorder (e.g., an "effective amount"). A proliferative disorder is "treated" when administration of reovirus to the proliferating cells effects lysis of the proliferating cells. This may result in a reduction in size of the neoplasm, or in a complete elimination of the neoplasm. The reduction in size of the neoplasm, or elimination of the neoplasm, is generally caused by lysis of neoplastic cells ("oncolysis") by the reovirus. Preferably the effective amount is that amount able
20 to inhibit tumor cell growth. Preferably the effective amount is from about 1.0 pfu/kg body weight to about 10^{15} pfu/kg body weight, more preferably from about 10^2 pfu/kg body weight to about 10^{13} pfu/kg body weight. For example, for treatment of a human, approximately 10^2 to 10^{17} plaque forming units (PFU) of reovirus can be used, depending on the type, size and number of tumors present.
25 The effective amount will be determined on an individual basis and may be based, at least in part, on consideration of the type of reovirus; the chosen route of administration; the individual's size, age, gender; the severity of the patient's symptoms; the size and other characteristics of the neoplasm; and the like. The

course of therapy may last from several days to several months or until diminution of the disease is achieved.

The immunosuppressant or immunoinhibitor and reovirus can be administered in a single dose, or multiple doses (i.e., more than one dose). The multiple doses can be administered concurrently, or consecutively (e.g., over a period of days or weeks). The reovirus can also be administered to more than one neoplasm in the same individual.

The compositions are preferably formulated in a unit dosage form, each dosage containing an appropriate amount of immunosuppressant or immunoinhibitor and from about 10^2 pfus to about 10^{13} pfus of the reovirus. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of reovirus calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

As mentioned above, it has been found that the reovirus is effective for the treatment of solid neoplasms in immunocompetent mammals. Administration of unmodified reovirus directly to the neoplasm results in oncolysis of the neoplastic cells and reduction in the size of the tumor in immunocompetent animals. When animals are rendered immunosuppressed or immunodeficient in some way, systemic administration of reovirus will be more effective in producing oncolysis.

It is contemplated that the reovirus may be administered in conjunction with surgery or removal of the neoplasm. Therefore, provided herewith are methods for the treatment of a solid neoplasm comprising surgical removal of the neoplasm and administration of a reovirus at or near to the site of the neoplasm.

It is contemplated that the reovirus may be administered in conjunction with or in addition to radiation therapy which renders the mammal immunosuppressed.

It is further contemplated that the reovirus of the present invention may be administered in conjunction with or in addition to known anticancer compounds or chemotherapeutic agents. Chemotherapeutic agents are compounds which may inhibit the growth of tumors. Such agents, include, but are not limited to, 5-fluorouracil, mitomycin C, methotrexate, hydroxyurea, cyclophosphamide, dacarbazine, mitoxantrone, anthracyclins (Epirubicin and Doxorubicin), antibodies to receptors, such as herceptin, etoposide, pregnasome, platinum compounds such as carboplatin and cisplatin, taxanes such as taxol and taxotere, hormone therapies such as tamoxifen and anti-estrogens, interferons, aromatase inhibitors, progestational agents and LHRH analogs.

The reovirus and immunosuppressants of the present invention have been found to reduce the growth of tumors that are metastatic. In an embodiment of the invention, a method is provided for reducing the growth of metastatic tumors in a mammal comprising administering an effective amount of a reovirus to the immunosuppressed mammal.

Utility

The diagnostic methods of the present invention may be used to identify the susceptibility of cells to reovirus infection by measuring constitutive ras-MAP signaling. This will be useful to determine in what cases reovirus treatment of cellular proliferative disorders is likely to be effective.

The reoviruses and immunosuppressants of the present invention may be used for a variety of purposes. They may be used in methods for treating proliferative disorders which exhibit constitutive MAPK phosphorylation in a mammal. They may be used to reduce or eliminate neoplasms. They may be used in methods for treating metastases. They may be used in conjunction with known treatments for cancer including surgery, chemotherapy and radiation.

In order to further illustrate the present invention and advantages thereof, the following specific examples are given but are not meant to limit the scope of the claims in any way.

EXAMPLES

In the examples below, all temperatures are in degrees Celsius (unless otherwise indicated) and all percentages are weight percentages (also unless otherwise indicated).

In the examples below, the following abbreviations have the following meanings. If an abbreviation is not defined, it has its generally accepted meaning:

μ M	=	micromolar
mM	=	millimolar
M	=	molar
ml	=	milliliter

	μ l	=	microliter
	mg	=	milligram
	μ g	=	microgram
	PAGE	=	polyacrylamide gel electrophoresis
5	rpm	=	revolutions per minute
	FBS	=	fetal bovine serum
	DTT	=	dithiothrietol
	SDS	=	sodium dodecyl sulfate
	PBS	=	phosphate buffered saline
10	DMEM	=	Dulbecco's modified Eagle's medium
	α -MEM	=	α -modified Eagle's medium
	β -ME	=	β -mercaptoethanol
	MOI	=	multiplicity of infection
	PFU	=	plaque forming units
15	MAPK	=	MAP kinase
	phosph-MAPK	=	phosphorylated-MAP kinase
	HRP	=	horseradish-peroxidase
	PKR	=	double-stranded RNA activated protein kinase
	RT-PCR	=	reverse transcriptase-polymerase chain reaction
20	GAPDH	=	glyceraldehyde-3-phosphate dehydrogenase
	EGFR	=	epidermal growth factor receptors
	MEK kinase	=	mitogen-activated extracellular signal-regulated kinase
	DMSO	=	dimethylsulfoxide
25	SCID	=	severe combined immunodeficiency

General Methods

Cells and Virus

Parental NIH-3T3 cells along with NIH-3T3 cells transformed with *v-Src* were a generous gift of Dr. Jove (University of Florida). Breast tumor cell lines

MDA-MB-468, MCF7, MDA-MB-435, T-47D, SK-BR-3 and the control HBL-100 cells, were a generous gift of Dr. Karl Riabowol (University of Calgary). All cell lines were grown in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS).

- 5 The Dearing strain of reovirus serotype 3 used in these studies was propagated in suspension cultures of L cells and purified according to Smith et al (1969) with the exception that β -mercaptoethanol (β -ME) was omitted from the extraction buffer.

Infection of Cells and Quantitation of Virus

- 10 Confluent monolayers of cells were grown in 24-well plates and infected with reovirus at an estimated multiplicity of infection of 80 PFU/cell. After 1 hr incubation at 37°C, the monolayers were washed with warm DMEM-10% FBS, and then incubated in the same medium. At various times postinfection, a mixture of NP-40 and sodium deoxycholate was added directly to the medium on the
15 infected monolayers to final concentrations of 1% and 0.5% respectively. The lysates were then harvested and virus yields were determined by plaque titration on L-929 cells.

Radiolabelling of reovirus-infected cells and preparation of lysates

- 20 Subconfluent monolayers (80% confluent) of cells were infected with reovirus (MOI~ 10 PFU/cell). At 46 hours after infection, the media was replaced with methionine-free DMEM containing 10% FBS and 0.1 mCi/ml [³⁵S]-methionine. After further incubation for 2 hours at 37°C, the cells were washed in phosphate-buffered saline (PBS) and lysed in the same buffer containing 1% Triton X-100, and 0.5% sodium deoxycholate. Lysates were then boiled and
25 stored at -70°C until use.

Immunoprecipitation and SDS-PAGE analysis

Immunoprecipitation of ^{35}S -labeled reovirus-infected cell lysates with anti-reovirus serotype 3 serum was carried out as previously described (Lee, P.W.K. *et al.* (1981) *Virology*, 108:134-146).

Detection of Phospho-MAPK and Total MAPK

- 5 The 'PhosphoPlus' p44/42 MAP kinase (Thr202/Tyr204) antibody kit (New England Biolabs) was used for the detection of MAP kinase in cell lysates according to the manufacturer's instructions. Briefly, subconfluent monolayer cultures were lysed with the recommended SDS-containing sample buffer, and subjected to SDS-PAGE, followed by electroblotting onto nitrocellulose paper.
- 10 The membrane was then probed with the primary antibody (anti-total MAPK or anti-phospho-MAPK), followed by the horseradish peroxidase (HRP)-conjugated secondary antibody as described in the manufacturer's manual.

Severe Combined Immune Deficient (SCID) Mice

- Five to eight week old male SCID mice were purchased from Charles River Canada and were treated according to a protocol approved by the University of Calgary Animal Care Committee.
- 15

Implantation of Allografts and Xenografts

- Actively growing MDA-MB-468 human breast cancer cells were harvested, washed, and resuspended in sterile PBS at a density of 2×10^7 cells/ml. 2.0×10^6 cells in 100 μl were injected subcutaneously at a site overlying the hind flank. Implanted tumors were allowed to grow for 2-3 weeks until palpable tumors of 0.5 x 0.5 cm were obtained.
- 20

intratumoral Injection of Reovirus

- Once the established tumors obtained a treatable size, a single intratumoral injection of 1.0×10^7 PFUs of either live or UV-inactivated reovirus (serotype 3, strain Dearing) in 20 μl sterile PBS. Tumor size was measured twice weekly for a
- 25

period of two to four weeks. Animals were sacrificed when they displayed severe morbidity due to excessive tumor burden or any observable level of distress.

Histoimmunological Analysis of Reovirus Infection

Immunofluorescent analysis was carried out using formalin-fixed, paraffin
5 embedded tumor sections mounted on cover slips. After removal of paraffin by
xylene, the sections were rehydrated, and exposed to the primary antibody (rabbit
polyclonal anti-reovirus type 3 serum diluted 1/100 in PBS) for 2 h at room
temperature. Following three washes with PBS, the sections were exposed to the
secondary antibody [goat anti-rabbit IgG (whole molecule)-fluorescein
10 isothiocyanate conjugate (FITC) or depending on the experiment, the same
concentration of Cy3, diluted 1/100 in PBS containing 10% goat serum and
0.005% Evan's Blue counterstain] for 1 h at room temperature. As an additional
counterstain, the nuclear stain DAPI, was also used. Finally, the fixed and treated
sections were washed three more times with PBS and then once with double-
15 distilled water. Slides were then dried and mounted on slides in 90% glycerol
containing 0.1% phenylenediamine, and viewed with a Zeiss Axiophot microscope
on which a Carl Zeiss camera was mounted (the magnification for all pictures was
200X).

Reovirus Infection of Primary Breast Tumor Samples

20 Biopsy breast tumor samples were sterilized by immersion in 95% ethanol
followed by several washes of sterile PBS. The sample was then sliced into small
sections and placed in a 24 well plate containing DMEM with 10% FCS.
Reovirus was added (1×10^8 PFUs). At various times post infection, samples were
washed in sterile PBS and then fixed in formalin. Samples were then embedded in
25 paraffin and sectioned for use in immunohistochemical analysis using antibodies
directed against total reovirus proteins.

Example 1. Transformation by the v-src Non-receptor Tyrosine Kinase Family Member Confers Susceptibility to Reovirus Infection

Although we have previously demonstrated that transfection of uninfected NR6 and NIH-3T3 cells with receptor tyrosine kinases was sufficient to allow reovirus replication (Strong, 1993), whether transfection of non-receptor tyrosine kinases could result in reovirus susceptibility was unknown. To determine whether activation of Src family kinases, which frequently contribute to the uncontrolled proliferation of many breast cancers, results in sufficient Ras activity to enable reovirus replication, uninfected NIH-3T3 cells were transformed with *v-src* and susceptibility to reovirus was assessed. Confluent monolayers of *v-src* transformed NIH-3T3 or their parental cells were exposed to reovirus at a multiplicity of infection (MOI) of ~50 plaque forming units (PFUs) of reovirus. Cells and media were harvested at various times after infection and resultant samples were used for plaque titration analysis to determine reovirus replication. Our results show dramatic cytopathic effect in the *v-src* transformed cells by 48 hours postinfection (data not shown) as well as an enhanced reovirus output as measured by plaque titration assay. These results (Figure 1) show that even non-receptor tyrosine kinases could potentially mediate reovirus infection and thus may be able to target an additional subset of breast tumors.

Example 2. Reovirus Can Infect a Panel of Human Breast Cancer Derived Cell Lines in Which Constitutive Ras/MAPK signaling Can Be Assessed

To assess the feasibility of using reovirus against breast derived tumors a panel of breast cancer cells including: MDA-MB-468, MCF7, MDA-MB-435, T-47D, and SK-BR-3 was selected and susceptibility to reovirus infection *in vitro* was determined. As a control, HBL-100 cells, which are derived from normal breast tissue were used. These six cell lines were grown to 80% confluency and were then challenged with reovirus at an MOI of 10. Cells were labeled with [³⁵S] methionine for a two-hour period at 48 hours postinfection. Cells were then washed in phosphate-buffered saline and were lysed. The prepared lysates were then used for immunoprecipitation using antibodies directed against total reovirus

protein. Immunoprecipitated proteins were analyzed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). The results (Figure 2A) clearly demonstrate effective reovirus replication within the cell lines derived from breast tumor, however reovirus replication was restricted in the HBL-100 cell line. This suggests that these breast tumor cells are infectable as a result of an activation of Ras, either through a direct mutation or via an upstream signaling element. It is significant in that it suggests that a high percentage of breast cancers could potentially be treatable by therapeutics that target Ras, although this specific activation is rare in this cancer type.

As MAPK phosphorylation is the inevitable result of signaling through Ras, the phosphorylated state of the protein should only be observed in a cell line with normal Ras signaling if a mitogenic stimulus is present. In a cell line with normal Ras signaling, the removal of these mitogenic stimuli should result in the cessation of signal through this pathway with a resultant dephosphorylation of MAPK. In a cell with aberrant Ras activity, either directly through mutational activation of Ras or alternatively via an upstream element, the phosphorylated form of MAPK should persist in either the presence or the absence of serum. To establish that the observed infectability in the breast tumor cell lines is a result of the activation of the Ras/MAPK signaling pathway, Western blot analysis of these cell lines was conducted using antibodies directed against phospho-MAPK. MDA-MB-468, MCF7, MDA-MB-435, T-47D, SK-BR-3 and HBL-100 cells were plated in a six well plate. Cells were then grown in 10% fetal calf serum (FCS) or were serum starved (0.5% FCS) for 48 hours. Cells were then washed in phosphate buffered saline (PBS) and were harvested in protein sample buffer. Cell lysates were subjected to SDS-PAGE and were subsequently blotted onto nitrocellulose paper and probed with anti-phospho-MAPK antibodies as recommended by the manufacturer. The results (Figure 2B) clearly demonstrate that the only uninfected cell line, HBL-100, possesses phospho-MAPK in the presence of serum and not in its absence, indicating that Ras signalling in this pathway is not aberrant. The remaining infected cell lines have phospho-MAPK independent of

the presence of mitogenic signals, as would be expected if they had constitutive activation of this pathway. Protein concentrations were standardized using antibodies directed against total MAPK (Figure 2C).

Example 3. Reovirus Can Act as an Anti-tumor Agent Against Breast Tumors *In Vivo*.

5 The human breast cancer cell line, MDA-MB-468 was introduced as tumor xenografts subcutaneously into a site overlying the hind flank of SCID mice. Following palpable tumor establishment, the MDA-MB-468 tumor was administered a single intratumoral injection of 1.0×10^7 plaque forming units
10 (PFUs) of reovirus serotype 3 (strain Dearing) in PBS. Control animals were given an intratumoral injection of UV-inactivated reovirus. Tumor growth was followed for a period of four weeks. As in demonstrated (Figure 3), the reovirus treatment results in a dramatic regression in tumor size. As before, hematoxylin/eosin (HE) staining of the remaining mass indicated that the single
15 reovirus injection results in the complete destruction of the tumor cells as compared to control tumors (data not shown). To determine if the observed lysis of the tumor cells was due to viral replication and to determine if there was spread of reovirus proteins beyond the tumor mass, immunofluorescence microscopy was conducted. Using antibodies directed against total reovirus proteins and paraffin
20 embedded thin sections of the remaining tumor mass it was determined that reovirus replication is restricted to the MDA-MB-468 tumor cells and does not extend into the surrounding skeletal muscle.

Example 4: Reovirus can Replicate in Primary Breast Specimens

To ensure that the oncolytic effect of reovirus is not due to an innate
25 characteristic of a passaged cell line, primary samples of human breast cancers were obtained to assess for reovirus infectability. Biopsy breast tumor samples were sterilized by immersion in 95% ethanol followed by several washes of sterile PBS. The sample was then sliced into small sections and placed in a 24 well plate containing DMEM with 10% FCS. Reovirus was added (1×10^8 PFUs). At

various times post infection, samples were washed in sterile PBS and then fixed in formalin. Samples were then embedded in paraffin and sectioned for use in immunohistochemical analysis using antibodies directed against total reovirus proteins. The results (not shown) clearly showed reovirus staining in those tumors
5 challenged, demonstrating viral replication in these primary samples.

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the invention as defined by the appended
10 claims.